

# Figures

1.1	Thesis structure.	21
3.1	Location of the study area in southern France.	36
3.2a	Trends in forest area and population in France since 1000 AD (Mather et al., 1999).	40
3.2b	Trends in forest area and rural population in France since 1840 AD (Mather et al., 1999).	40
4.1	Hierarchical land cover classification scheme based on Tomaselli (1981).	46
4.2	Change classes used for visual interpretation. % Cover expressed as the % of the total study area.	47
4.3	Change class cross tabulations for the continuous variables. The key of the change classes is given in figure 4.2.	49
4.4	Cross tabulations between the change class and lithology maps.	50
4.5	Percentages of the five most important soil classes of abandoned lands as a function of the time since abandonment.	52
4.6	Important soil classes per land cover change class and the most important vegetation classes per soil type for change class 1: recently abandoned fields. % Refers to the % cover in the study area.	52
4.7	Important soil classes per land cover change class and the most important vegetation classes per soil type for the areas with observed change and that are abandoned before the 1940's (change class 2 to 6). % refers to the % cover in the study area.	53
4.8	Important soil classes for (change class 9) and the most important vegetation classes per soil type for the unchanged natural areas, interpreted as the climax stage of the vegetation. % Refers to the % cover in the study area.	54
4.9	Transition paths per Time Since Abandonment (TSA) class for the recent abandoned fields (change class 1). % Refers to the % cover in the study area.	55
5.1	Hierarchical land cover classification scheme, modified from Tomaselli (1981).	61
6.1	Simulated 3 x 3 kernels of different natural vegetation types with corresponding adjacency event matrix. Only the upper triangular is considered because $M_{ij} = M_{ji}$ . (Barnsley and Barr, 1996).	71
6.2	Hierarchical land cover classification scheme used for field survey and for image analysis based on Tomaselli (1981). Classes used in the 12-class SPARK classification are shown in bold.	74
6.3	Maximum likelihood overall accuracy values for 7, 10, 12 and 15 classes.	76
7.1	Hierarchical land cover classification scheme, modified from Tomaselli (1981).	84
8.1	Hierarchical land cover classification scheme, modified from Tomaselli (1981).	100
A1	HyMap image (R,G,B = band 29, band 15, band 9).	114
A2	ASTER image (R,G,B = band 3, band 2, band 1).	115
A3	Landsat 7 ETM+ image (R,G,B = band 4, band 3, band 2).	115

A <sub>4</sub>	Aerial photograph 1946.	116
A <sub>5</sub>	Aerial photograph 1970.	116
A <sub>6</sub>	Aerial photograph 2001.	117
A <sub>7a, b</sub>	Spatial distribution of change classes in the study area. Vailhan area on the left, Mourèze area on the right.	118
A <sub>8</sub>	15 Class Random Forests (RF) classification result with inclusion of ancillary data (scenario 4, table 5.2).	119
A <sub>9a</sub>	DAIS7915 colour composite (R,G,B = PCA <sub>1</sub> ,PCA <sub>2</sub> ,PCA <sub>3</sub> ). Green corresponds to dense vegetation, red to bare soil and agriculture.	120
A <sub>9b</sub>	Maximum likelihood classification result with 7 classes: the heterogeneous vegetation in the image is classified in homogeneous clumps of trees, shrubs and grass.	120
A <sub>10a</sub>	Initial NDVI-land cover map with 27 NDVI-classes.	121
A <sub>10b</sub>	12-Class maximum likelihood classification results.	121
A <sub>10c</sub>	SPARK classification results with kernel size 3 and 27 NDVI-classes.	122
A <sub>10d</sub>	SPARK classification results with kernel size 11 and 10 NDVI-classes.	122
A <sub>11</sub>	Upper: subset of the 12-Class maximum likelihood classification results (overall accuracy: 56.2%). Lower: the SPARK classification results with kernel size 3 and 27 NDVI-classes (overall accuracy 64.5%). The red line roughly indicates the border between the maquis dominated vegetation types in the north and the garrigue dominated vegetation types in the south.	123
A <sub>12</sub>	Soil map of the study area and SAM rule image for dense matorral (class 2).	124
A <sub>13</sub>	18 Class MLH result (a,b) and 18 class ADCM result (c,d). Numbers in the map refer to the four sub-areas of figure A <sub>14</sub> .	125
A <sub>14</sub>	18 Class MLH result (left) and 18 class ADCM result (right) in four sub-areas. Area numbers refer to the areas indicated in figure A <sub>13</sub> .	126
A <sub>15</sub>	Model results for time step 1, 1946 (A, B), time step 57, 2003 (C, D), time step 100, 2046 (E, F) and classification results of the HyMap 2003 image (G, H). Numbers in the map refer to the four sub-areas (continued on next page).	127